

F-Distribution

Explorable.com 47.7K reads

The F-distribution, also known as the Snedecor's F-distribution or the Fisher-Snedecor distribution (after R.A. Fisher and George W. Snedecor), is the distribution of ratios of two independent estimators of the population variances.

Suppose we have two samples with n_1 and n_2 observations, the ratio $F = s_1^2 / s_2^2$ where s_1^2 and s_2^2 are the sample variances, is distributed according to an F-distribution with $v_1 = n_1 - 1$ numerator degrees of freedom, and $v_2 = n_2 - 1$ denominator degrees of freedom.

For example, if F follows an F-distribution and the degrees of freedom for the numerator are 4 and the degrees of freedom for the denominator are 10, then $F \sim F_{4,10}$. For each combination of these degrees of freedom there is a different F-distribution. The F-distribution is most spread out when the degrees of freedom are small. As the degrees of freedom increase, the F-distribution is less dispersed.



The banner features the Explorable logo (a flask icon) and the text "EXPLORABLE Quiz Time!". Below the logo are three quiz cards:

- Quiz: Psychology 101 Part 2 (with an image of roller skates)
- Quiz: Psychology 101 Part 2 (with an image of colored pencils)
- Quiz: Flags in Europe (with an image of a Ferris wheel)

A link "See all quizzes =>" is located at the bottom right of the banner.

Properties

The F-distribution has the following properties:

The mean of the distribution is equal to $v_1 / (v_2 - 2)$. The variance is equal to $[v_2^2 * (v_1 + 2)] / [v_1 * (v_2 - 2) * (v_2 - 4)]$

The F-distribution is skewed to the right, and the F-values can be only positive. The curve reaches a peak not far to the right of 0, and then gradually approaches the horizontal axis. The F-distribution approaches, but never quite touches the horizontal axis.

Uses

The main use of F-distribution is to test whether two independent [samples](#) [1] have been drawn for the [normal populations](#) [2] with the same variance, or if two independent estimates of the population variance are homogeneous or not, since it is often desirable to compare two [variances](#) [3] rather than two [averages](#) [4]. For instance, college administrators would prefer two college professors grading exams to have the same variation in their grading. For this, the [F-test](#) [5] can be used, and after examining the [p-value](#) [6], inference can be drawn on the variation.

Assumptions

In order to perform F-test of two variances, it is important that the following are true:

- The [populations](#) [7] from which the two samples are drawn are [normally distributed](#) [2].
- The two populations are independent of each other.

If the two populations have equal variances, then s_1^2 and s_2^2 are close in value and F is close to 1. But if the two population variances are very different, s_1^2 and s_2^2 tend to be very different, too.

Choosing s_1^2 as the larger sample variance causes the ratio to be greater than 1. If s_1^2 and s_2^2 are far apart, then F is a large number. Therefore, if F is close to 1, the evidence favours the [null hypothesis](#) [8] (the two population variances are equal). But if F is much larger than 1, then the [evidence](#) [9] is against the null hypothesis, and we can infer that possibly the population variances differ to a large extent.

Anova and F

In the technique known as [Analysis of Variance](#) [10] (ANOVA) which plays a very important role in [Design of Experiments](#) [11], the variance ratio test is applied to [test the significance](#) [12] of different components of variation against error variation.

For example, a new drug for treating Osteoporosis could need to be field tested. Since severity of this disease is generally a function of age, the new drug could be administered randomly to n patients in each age group. Put differently, this would be an experiment in m age groups and n different dosage levels of the drug allocated randomly to the patients. With figures provided from patients for each age group \times dose combination, we can use the variance ratio test ([F-test](#) [5]) to test for difference between dose levels and if this variation can be attributed to chance.

The other uses include testing the significance of the [correlation](#) [13] ratio between two random variables, and to test the [linearity of regression](#) [14].

Source URL: <https://forum.explorable.com/f-distribution>

Links

- [1] <https://forum.explorable.com/sample-group>
- [2] <https://forum.explorable.com/normal-probability-distribution>
- [3] <https://forum.explorable.com/statistical-variance>
- [4] <https://forum.explorable.com/arithmatic-mean>
- [5] <https://forum.explorable.com/f-test>
- [6] <https://forum.explorable.com/p-value>
- [7] <https://forum.explorable.com/population-sampling>
- [8] <https://forum.explorable.com/null-hypothesis>

- [9] <https://forum.explorable.com/empirical-evidence>
- [10] <https://forum.explorable.com/anova>
- [11] <https://forum.explorable.com/design-of-experiment>
- [12] <https://forum.explorable.com/significance-test>
- [13] <https://forum.explorable.com/statistical-correlation>
- [14] <https://forum.explorable.com/linear-regression-analysis>